

Lorentz Covariant Spin–Grouping of Baryon Resonances

M. Kirchbach *

Institute for Nuclear Physics, University Mainz, D-55099 Mainz, Germany

A well pronounced spin–grouping of baryon resonances to $O(4)$ partial waves is found in baryon spectra and shown to be well interpreted in terms of Lorentz group representations of the type $\{\frac{1}{2} + l', \frac{1}{2} + l'\} \otimes [\{\frac{1}{2}, 0\} \oplus \{0, \frac{1}{2}\}]$ with l' integer. In this way the relativistic description of finite dimensional resonance towers containing higher-spin states becomes possible. It is further argued that the nucleon excitations into the $l' = 1$ and $l' = 2$ multiplets are chiral phase transitions.

The relativistic description of resonances with higher-spins is one of the oldest unsolved problems of baryon spectroscopy. Attempts to handle it in the mid to late 60's failed through the improper choice for the irreducible representations of the Lorentz group considered to embed the baryon excitations known at that time. For example, Kleinert considered in [1] unitary representations of the Lorentz group describing a spin-tower of states without upper end starting for the nucleon $P_{11}(939)$ with the $D_{13}(1520)$, $F_{15}(1680)$, $G_{17}(2190)$ etc. resonances. Alternatively, Regge phenomenology required in the t-channel near origin (i.e. $t_\mu \approx 0$) a grouping of approximately mass degenerate baryon Regge poles into the finite dimensional non-unitary representations of the Lorentz group of the type $\{l_1, l_2 \pm \frac{1}{2}\}$ with integer l_i 's (so called Toller poles [2]). Unfortunately, both the schemes mentioned above are in disagreement with data, in particular because the parity duplication of the resonances required by them is absent from the spectra. On the other side, Weinberg described baryon resonances as isolated $\{J, 0\} \oplus \{0, J\}$ higher–spin states. In order to obtain the corresponding relativistic equations of motion Weinberg put forward the idea in [3] to embed the $\{J, 0\} \oplus \{0, J\}$ state, considered as pointlike, within the Rarita–Schwinger multiplet $\{\frac{1}{2}k, \frac{1}{2}k\} \otimes [\{\frac{1}{2}, 0\} \oplus \{0, \frac{1}{2}\}]$ (with k integer and related to J via $k = J - 1/2$) and to eliminate all the lower–spin components, considered as *redundant unphysical states* by means of a set of suitably chosen auxiliary conditions. Within Weinberg's scheme a spin- J resonance of mass M and momentum p_μ is described by a totally symmetric traceless rank- k Lorentz tensor with Dirac spinor components $\Psi_{\mu\mu_1\dots\mu_{k-1}}$ satisfying the conditions

$$(\gamma \cdot p - M) \Psi_{\mu\mu_1\dots\mu_{k-1}} = 0, \quad (1)$$

$$p^\mu \Psi_{\mu\mu_1\dots\mu_{k-1}} = 0, \quad \gamma^\mu \Psi_{\mu\mu_1\dots\mu_{k-1}} = 0. \quad (2)$$

Here, Proca's equation in Eq. (2) eliminates the spin-0 and preserves the spin-1 components from the $\{1/2, 1/2\}$ representations as associated with the Lorentz indices. The second set of auxiliary conditions in (2) eliminates then the longitudinal components of the spin-1 field. It ensures that the state surviving the elimination procedure carries the maximal allowed helicities and describes, therefore, the highest–spin state of the Rarita–Schwinger multiplet. A serious disadvantage of the Rarita–Schwinger scheme for treating higher–spin states is that for off–shell particles the auxiliary conditions are not any longer valid and the appearance of arbitrary unphysical parameters in the Lagrangian describing the couplings of the states under consideration to external fields is inevitable. Because of all these difficulties, the requirement for relativistic and thereby for field theoretical description of higher–spin resonances was gradually given up by hadron spectroscopy thus opening space for pragmatic evidently non–relativistic schemes such as the $O(3)_L \otimes SU(6)_{SF}$ hadron book–keeping symmetry. Through this group the trivial spin-flavor (SF) correlation between three quarks in the 1s-shell was naively extended to arbitrary orbital angular momenta. The problems raised by symmetry groups of strong interaction based on tight correlations between the spin- and flavor degrees of freedom of quarks not only have conceptual disadvantages, but also their predictions are not satisfactory. For example, the $O(3)_L \otimes SU(6)_{SF}$ classification scheme predicts a substantial excess of baryon excitations known as 'missing resonances'. Finally, the basic idea of the multiplets as well separated families of particles of different internal but same space–time properties, seems quite inappropriate here, where the spacing between the $O(3)_L \otimes SU(6)_{SF}$ multiplets is much smaller as compared to the mass splitting within the multiplets.

On the other side, speed plot analysis of the pole positions on the complex energy plane of various baryon resonances ($L_{2I,2J}$) with masses below ~ 2500 MeV performed by Höhler [4] revealed a well-pronounced spin–clustering in baryon spectra. This is quite a surprising result as it was not anticipated by any model or theory. Independently, the symmetry of all reported N , Δ and Λ baryon excitations with masses below 2500 MeV was re-analyzed in our previous work [5] and shown to be governed by $O(1,3)_{LS} \otimes SU(2)_I$ rather than by $O(3)_L \otimes SU(6)_{SF}$ as predominantly used since the invention of the naive three flavor quark model. The $O(1,3)_{LS} \otimes SU(2)_I$ symmetry indicates that

*E-mail address: mariana@kph.uni-mainz.de

the spin-orbital correlation between quarks is much stronger than the spin-flavor one. Indeed, it was demonstrated in [5] that Höhler's poles correspond to $O(4)$ partial waves as they are identical to Lorentz multiplets of the type $\{\frac{1}{2} + l', \frac{1}{2} + l'\} \otimes [\{1/2, 0\} \oplus \{0, 1/2\}]$ with l' integer. All orbital angular momenta (denoted here by l) contained within a $\{\frac{1}{2} + l', \frac{1}{2} + l'\}$ multiplet have either natural or unnatural parity¹. Coupling a Dirac particle to l is then standard and leads to covariantly transforming spin-groups of states with $\vec{J} = \vec{l} \otimes \frac{\vec{1}}{2}$ and $l = 0, \dots, 2l' + 1$ all having either natural or unnatural parities according to $J = \frac{1}{2}^+, \frac{1}{2}^-, \frac{3}{2}^-, \dots, (k + \frac{1}{2})^\pi$ with $k = 2l' + 1$, $\pi = (-1)^l$ for natural, and $\pi = (-1)^{l+1}$ for unnatural parities. Note that the $\{\frac{1}{2} + l', \frac{1}{2} + l'\}$ multiplets are well known from the Coulomb problem, where they correspond to even principal quantum numbers $n = k + 1$. These finite dimensional unitary representations of $O(4)$, the maximal compact group of the Lorentz group, contain only once each $O(3)$ representation corresponding to an intrinsic orbital angular momentum $l = 0, \dots, k$. The Coulomb multiplets yield in the $O(1,3)$ group finite dimensional *non-unitary* representations. Because of the possibility of restricting $O(1,3)$ to $SU(2) \otimes SU(2)$ for the $\{\frac{1}{2} + l', \frac{1}{2} + l'\}$ representations, however, the latter may be used for the construction of relativistic equations of motion. The baryon resonances are now collected into the following representations [5]:

$$\begin{aligned}
l' = 0 : \quad \Psi_\mu : P_{2I,1}; S_{2I,1}, D_{2I,3}, \quad \text{for } I = 0, \frac{1}{2}, \frac{3}{2}, \quad \text{and} \\
l' = 1 : \quad \Psi_{\mu\mu_1\mu_2} : S_{2I,1}; P_{2I,1}P_{2I,3}; D_{2I,3}, D_{2I,5}; F_{2I,5}, F_{2I,7}, \\
l' = 2 : \quad \Psi_{\mu\mu_1\dots\mu_4} : S_{2I,1}; P_{2I,1}P_{2I,3}; D_{2I,3}, D_{2I,5}; F_{2I,5}, F_{2I,7}; \\
G_{2I,7}, G_{2I,9}; H_{2I,9}, H_{2I,11}, \quad \text{for } I = \frac{1}{2}, \frac{3}{2}. \tag{3}
\end{aligned}$$

Note that the states with the highest spins $J^\pi = (k + \frac{1}{2})^\pi$ from the Lorentz multiplets under consideration and the corresponding isolated $\{J^\pi, 0\} \oplus \{0, J^\pi\}$ states may belong to different Fock spaces due to different origins for their parities. Consider, as an illustration, a spin- $3/2^-$ resonance to emerge as the highest spin state of the Lorentz multiplet $\{1/2, 1/2\} \otimes [\{1/2, 0\} \oplus \{0, 1/2\}]$. The negative parity of this resonance signals an internal P wave state. Such a natural parity state resides within a Fock space built upon a scalar vacuum. On the contrary, the negative parity of an isolated $\{3/2^-, 0\} \oplus \{0, 3/2^-\}$ state arises necessarily from the coupling of the fermion degree of freedom, when considered as fundamental, to an internal 0^- state. Such an unnatural parity state necessarily resides within a Fock space built upon a pseudoscalar vacuum. This aspect of mapping isolated higher-spin states onto multi-spinor representations of the Lorentz group is of crucial importance for selecting baryons as genuine parity partners and, therefore, for establishing the scale of chiral symmetry restoration (see below).

Considering resonances as approximate $O(4)$ partial waves brings numerous advantages relative treating them as $O(3)$ partial waves. First of all, the number of the ‘missing resonances’ is strongly reduced. Indeed, within that scheme the Δ spectrum below 2000 MeV appears complete as all observed states fit into the $l' = 0$, and $l' = 1$ Lorentz multiplets in Eq. (2). As the F_{37} state from the $l' = 1$ Lorentz multiplet has to be paralleled in the nucleon sector by a (not reported) F_{17} resonance with a mass around 1700 MeV, only that latter state has to be viewed as a ‘missing resonance’ among the non-strange baryon excitations with masses below 2000 MeV. In continuing by comparing the states from the third nucleon and Δ clusters with $l'=2$, four more ‘missing resonances’ are predicted. These are the $H_{1,11}$, P_{31} , P_{33} , and D_{33} states with masses between 2200 and 2400 MeV. In summary, five new, still unobserved non-strange resonances have been predicted in [5]. Remarkably, the spacing of about 200 MeV among the relativistic multiplets with masses below 2000 MeV appears now much larger as compared to the maximal mass splitting of 50-70 MeV between the corresponding multiplet members (see Fig. 1). The baryon classification scheme in Eq. (3) shows that the parities of the resonances from the first spin-clusters with $l' = 0$ are always natural. This means that they can be considered to emerge within a Fock space built upon a vacuum of positive parity as spontaneously selected in the course of the realization of chiral symmetry in the hidden Nambu–Goldstone mode in the low energy regime.

¹The parity $(-1)^{L+1}$ of a single πN resonance $L_{2I,2J}$ in standard notation [8], where L stands for the relative angular momentum in the decay channel, is determined in the present classification scheme by either $(-1)^l$ or $(-1)^{l+1}$, depending on whether the parity of the intrinsic orbital angular momentum is natural or unnatural. In the present notation, L takes the values of either $L = |l - 1|, (l + 1)$ for natural, or $L = l$ for unnatural parities.




Fig. 1 Baryon spectra in terms of the $\{j, j\} \otimes [\{\frac{1}{2}, 0\} \oplus \{0, \frac{1}{2}\}]$ representations. Only the $\{j, j\}$ assignments are given. Each dashed area contains $4j + 1$ states.

On the contrary, the resonances of the second and third spin-clusters with $l' = 1$, and $l' = 2$, respectively, have always unnatural parities and can be considered to reside within a Fock space built upon a vacuum of negative parity. Both the $\psi_N \rightarrow \Psi_{\mu\mu_1\mu_2}$, and $\psi_N \rightarrow \Psi_{\mu\mu_1\dots\mu_4}$ transitions are, therefore, in contrast to the $\psi_N \rightarrow \Psi_\mu$ excitation, chiral phase transitions. A further advantage of the $O(1,3)_{LS} \otimes SU(2)_I$ classification scheme for baryons is that it opens a new possibility for the relativistic description of the higher-spin resonances as members of covariant representations. Indeed, the relativistic propagators of the spin-clusters can be directly read off from the representation theory of the Lorentz group as

$$S_{\mu\mu_1\dots\mu_{k-1}; \nu\nu_1\dots\nu_{k-1}} = \frac{(\gamma \cdot p + M) \otimes_{n=1}^{n=k} (g_{\mu_n\nu_n} - \frac{1}{M^2} p_{\mu_n} p_{\nu_n})}{2M(p^2 - M^2)}, \quad (4)$$

with M standing for the mass of the degenerate resonances. For example, the $(S_{2I,1}, D_{2I,3})$ cluster is described in terms of the Lorentz vector with Dirac spinor components Ψ_μ from Eq. (3) and its propagator is given by [5]

$$S_{\mu\nu} = \frac{(\gamma \cdot p + M)(g_{\mu\nu} - \frac{1}{M^2} p_\mu p_\nu)}{2M(p^2 - M^2)}. \quad (5)$$

In noting that, say, the first S_{11} and D_{13} states appear separated by only 15 MeV, one sees that the relativistic contribution of these states to the amplitude of processes like meson photoproduction at threshold, can easily be calculated. Along the line of the representation theory of the Lorentz group, the construction of higher-cluster propagators and interactions with external fields is straightforward. For example, for the case of a $\mathcal{B} \rightarrow N + V$ process, where $\mathcal{B}$ stands for a Lorentz covariant spin-cluster, while V is a vector meson, a possible Lagrangian can be written as

$$\mathcal{L}_{\mathcal{B}VN} = \bar{\Psi}^{\mu\mu_1\dots\mu_{k-1}} \left(\frac{f_k}{m_\pi^{k-1}} \partial_{\mu_1} \dots \partial_{\mu_{k-1}} A_\mu + \frac{f'_k}{m_\pi^k} \partial_{\mu} \dots \partial_{\mu_{k-1}} A \right) \psi_N, \quad (6)$$

where $k = 2l' + 1$, A_μ denotes the vector meson field, while the coupling strengths f_k and f'_k can be extracted from data.

The Lorentz covariant spin-clusters of baryon resonances introduced above are in fact nothing else but the special case of the Rarita-Schwinger representations with odd k 's, the essential difference being that all the lower-spin states are no longer redundant components that need be eliminated, but *physically observable resonances* related to the composite character of the baryons. The Lorentz-Dirac index notation for the spin-clusters introduced in Eq. (3) indicates that the first resonance-tower will predominantly couple to the pion-nucleon (or η -nucleon) system carrying each one Dirac and one Lorentz index. The second and third spin-clusters will prefer couplings to multipion-nucleon final states (one Dirac- and several Lorentz indices) in agreement with the empirical observations. According to that, the reason for the suppression of the $S_{11}(1650) \rightarrow N + \eta$ decay channel as compared to the $S_{11}(1535) \rightarrow N + \eta$ one, can be a simple re-distribution of decay strength in favor of the new opened $S_{11}(1650) \rightarrow N + \pi + \pi$ channel in accordance with data. Now, the existence of the Lorentz covariant spin-grouping of baryon resonances can be interpreted to

emerge via the coupling of a quark to approximately mass-degenerated Di-quarks of spins $l = 0, \dots, 2l' + 1$ having all either natural or unnatural parities. Indeed, the covariant quark-Di-quark model based on solving the Bethe-Salpeter equation reveals an internal $O(4)$ symmetry [7] as visible from the rapid convergence of its solutions in the basis of the orthogonal (Gegenbauer) polynomials of the group $O(4)$ corresponding to four dimensional partial waves.

Thus the major advantage of our new relativistic spectrum generating algebra for baryons is that it reconciles such seemingly contrary ideas of the baryon dynamics like the constituent quark model on the one side, and the multi-spinor representations of the Lorentz group corresponding to structureless (pointlike) particles, on the other side. Now the apparent analogy between the spectrum of the hydrogen atom and the baryon spectra leads to the question whether the positions of the Lorentz covariant spin-clusters is determined by the inverse squared $1/n^2$ of the principle quantum number and follow a sort of modified Balmer-series pattern. The answer to that question is positive [5]. Below quite a simple empirical recursive relation is suggested that describes with quite an amazing accuracy the reported mass averages of the resonances from the Lorentz multiplets only in terms of the cluster quantum numbers and the masses of the ground state baryons:

$$M_n - M_{n-2} = \left(\frac{1}{(n-1)^2} - \frac{1}{n^2} \right) J_{n-2}^{max} (J_{n-2}^{max} + 1) M_{n-2},$$

$$n = 2l' + 2, \quad l' > 0, \quad J_{n-2}^{max} = n - \frac{1}{2}. \quad (7)$$

Here, M_n denotes the mass of the respective Lorentz multiplet. The position of the first excited nucleon spin-cluster with $l' = 0$ is given instead by (7) rather by $M_{n=2} - M_{n=1} = (\frac{1}{1^2} - \frac{1}{4})(1^2 - \frac{1}{4})M_{n=1}$ with $M_{n=1} = M_N$. For the Δ and Λ baryons, the mass scale $M_{n=1}$, has to be replaced by $(M_N + M_\Delta)/2$, and $(M_N + M_\Lambda)/2$, respectively (compare Table 2 of Ref. [6] for details). To conclude, baryon resonances group to $O(4)$ rather than to $O(3)$ partial waves, and the baryon spectra are evidently generated by the relativistic $o(1,3)_{LS} \otimes su(2)_I$ group algebra rather than by the algebra of the non-relativistic $O(3)_L \otimes SU(6)_{SF}$ group

Work supported by the Deutsche Forschungsgemeinschaft (SFB 201).

-
- [1] H. M. Kleinert, Fortschritte der Physik, **16**, 1 (1968).
 - [2] M. Toller, Nuovo Cim. **54A**, 295 (1968).
 - [3] S. Weinberg, Phys. Rev. **133**, B1318 (1964).
 - [4] G. Höhler, In: *Physics with GeV- Particle Beams* eds. H. Machner and K. Sistemich (World Scientific, Singapore) 1995, p. 198.
 - [5] M. Kirchbach, Mod. Phys. Lett. **A12**, 2373 (1997); *ibid.* 3177.
 - [6] M. Kirchbach, Mod. Phys. Lett. **A13**, 823 (1998).
 - [7] K. Kusaka *et al.*, Phys. Rev. **D55**, 5299 (1997).
 - [8] Particle Data Group, R. M. Barret *et al.*, Phys. Rev. **D54**, 1 (1996).